

# NALOXONE REFILL QUESTIONNAIRE

**FOR INTERVIEWER:** PLEASE ASK EACH QUESTION IN ORDER. CIRCLE THE RESPONSE THAT BEST MATCHES THE ANSWER GIVEN BY THE PARTICIPANT.

Date of interview: \_\_\_/\_\_\_/\_\_\_      Organization: \_\_\_\_\_ Staff Initials: \_\_\_\_\_

## A. INTRODUCTORY QUESTIONS

<b>1. What is your birth date?</b> Day _____ Month _____ Year _____					
<b>2. What is your gender?</b>					
01	Male	02	Female	03	Transgender/transsexual
<b>3. What is the highest level of education or schooling you have completed?</b>					
01	No schooling completed	03	High school graduate	05	College graduate
02	Incomplete high school	04	Some college or technical training	06	Graduate school
<b>4. What happened to the last dose of naloxone you were given?</b>					
01	Lost or stolen	03	It was confiscated by authorities	05	I used it during an overdose <i>[Skip to Section B]</i>
02	It broke	04	I gave it away or sold it	06	Other
<i>[To interviewer: please provide any relevant notes here]</i>					
<i>If the person answered 01, 02, 03, 04 or 06, thank them, provide them with a short refresher training, and give them new naloxone. If they answered 05, please continue to Section B.</i>					

## B. RECENT EXPERIENCE OF OVERDOSE

**Please answer these questions about the situation where the ampoule of naloxone was used.**

<b>5a. What is your relationship to the person that overdosed?</b>		01	Acquaintance	03	Boy/girlfriend or spouse
		02	Friend	04	Myself
		03	Stranger		
<i>If the person answers 04, skip to question 10d.</i>					
<b>5b. What was the gender of the person who overdosed?</b>					
01	Male	02	Female	03	Transgender/transsexual
<b>5c. What was the age of the person who overdosed?</b>					
01	Under 18	03	31-40	05	51 and above
02	18-30	04	41-50	06	Don't know
<b>5d. When did the overdose happen?</b>		Day _____ Month _____ Year _____			
<b>5e. What drugs were being used when the overdose happened?</b>		01	benzodiazepines (midazolam, xen, etc.)	06	other drugs (please indicate)
<i>To interviewer: please, circle around all that apply.</i>		02	opiates	07	Don't know/not sure
		03	stimulants (meth/amphetamines)		
		04	methadone		

	05	alcohol		
--	----	---------	--	--

<b>5f. Where were drugs being used when the overdose happened?</b>	01	In someone's home	04	other (please indicate)
	02	In the street or public place		_____
	03	Shooting gallery	05	Don't know/not sure

**5g. What is the closest intersection or address to the spot where the overdose took place?**  
 Intersection of street \_\_\_\_\_ and street \_\_\_\_\_

<b>5h. Who else was around the person experiencing an overdose?</b> <i>To interviewer: please, circle around all that apply.</i>	01	Other drug users	04	Non-drug-using friends
	02	Family members of the person overdosing	05	Strangers
	03	The victim's spouse or boy/girlfriend	06	No one
			07	Don't know

<b>6a. Had the person been released from prison or jail in the 6 months prior to the overdose?</b>	(0) no	(1) yes	(2) don't know
--	--------	---------	----------------

<b>6b. Had the person been released from an 06 center in the 6 months prior to the overdose?</b>	(0) no	(1) yes	(2) don't know
--	--------	---------	----------------

**7. During this overdose incident, what things did you or others do to try to help the victim (other than administering naloxone)?**  
*To interviewer: please, circle around all that apply.*

01	Called ambulance	06	Injected with stimulants, water or salt
02	Took to hospital	07	Called their name to try to wake them
03	Placed in the rescue position (on the side)	08	Did nothing
04	Performed rescue breathing/ heart massage/CPR	09	Other (specify): _____
05	Tried to wake person by causing pain, shower/bath, applying ice to wake the person, slapping or kicking person's chest		

**8a. How was the naloxone injected into the person experiencing an overdose?**

01	Into a muscle	02	Into a vein	03	Subcutaneously (under the skin)	04	Don't know
----	---------------	----	-------------	----	---------------------------------	----	------------

<b>8b. What type of needle was used to administer naloxone?</b>	(0) Intramuscular needle (21 or 23 gauge, usually 1 ½ inches or longer)	(1) intravenous needle (diabetic or tubercular syringe or similar; 25 gauge or higher)	(2) Don't know
---	---	--	----------------

<b>9a. After naloxone was given, did the overdose come back?</b>	(0) no	(1) yes	(2) don't know
--	--------	---------	----------------

<b>9b. If YES, did you give the person a second dose of naloxone?</b>	(0) no	(1) yes	(2) don't know
---	--------	---------	----------------

<b>9c. If YES, how long after the first naloxone dose was the 2<sup>nd</sup> one given?</b>	(0) less than 5	(1) more than 5	(2) don't know
---	-----------------	-----------------	----------------

			minutes	minutes	
<b>9d. How long after revival did someone stay with the overdose victim to make sure they were okay?</b>					
01	Left right away	02	Stayed for less than 2 hours	03	Stayed for 2 hours or more
<b>10. Did the person overdosing have any physical, emotional or social problems after the overdose? If so, please describe. To interviewer: please, circle around all that apply.</b>					
01	Withdrawal symptoms (nausea, achiness, vomiting)	02	Seizures	03	Pulmonary edema
04	Depression	05	Legal problems	06	Financial problems
		07	Guilt	08	Other emotional problems
		09	Other (specify): _____		
<b>11. Was the person taken to the hospital?</b>					
			(0) no	(1) yes	(2) Don't know
<b>12. Did the person who overdosed live or die?</b>					
			(0) survived	(1) died	(2) don't know
<b>13. Today, how confident are you that you can correctly use naloxone to reverse future overdoses?</b>					
01	Not at all confident	02	Hardly confident	03	Moderately confident
04	Completely confident				
<b>14. Today, how confident are you that you can do things to improve your own health?</b>					
01	Not at all confident	02	Hardly confident	03	Moderately confident
04	Completely confident				
<b>15. Today, how confident are you that you can improve the health of others?</b>					
01	Not at all confident	02	Hardly confident	03	Moderately confident
04	Completely confident				

**Thank the person for all the information they have provided. Give them a quick refresher training and two new ampoules of naloxone to take home.**